

Mean differences (in days) between exfoliation of the deciduous precursor and first appearance of a dental element of the permanent dentition for all breed sizes, calculated on exact exfoliation and emergence times of n number of teeth.

Tooth	All sizes			Giant			Large			Medium			Small			Toy		
	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD
Maxilla																		
I1	1109	0.83	1.56	112	0.68	1.41	347	1.01	1.64	210	1.26	1.02	309	0.26	1.53	131	2.1	4.17
I2	1068	0.65	1.3	120	1.02	2.54	332	0.3	0.87	182	0.93	1.15	310	0.68	0.8	124	0.82	1.78
I3	1061	0.53	0.6	124	0.12	0.42	336	0.53	0.54	196	0.78	0.53	283	0.69	0.75	122	0.05	0.19
C	741	-2.46	4.7	85	-5.13	3.68	219	-2.68	4.64	147	-0.3	3.19	210	-2.52	5.89	80	-3.84	2.97
P2	775	0.33	1.39	97	-0.69	1.87	227	0.31	1.26	132	0.41	0.41	223	-0.13	4.18	96	-0.34	1.14
P3	861	1.36	1.48	120	1.32	1.25	267	1.17	1.78	141	2.08	1.51	232	1.62	1.51	101	0.93	0.68
P4	879	0.66	1.61	113	0.51	0.59	270	1.02	1.42	168	1.09	0.73	236	0.05	2.39	92	0.18	0.65
Mandibula																		
I1	1074	0.82	1.16	119	0.42	0.48	338	0.62	0.75	189	0.99	0.65	308	1.73	2.27	120	-0.02	0.69
I2	1047	0.61	1.09	121	0.13	0.42	318	0.28	0.63	180	0.75	0.56	294	1.22	1.85	134	0.75	0.82
I3	1045	0.47	1.09	121	-0.68	2.13	322	0.58	0.75	188	0.83	0.55	279	0.5	0.93	135	0.79	0.89
C	742	-0.99	3.51	94	-1.46	2.16	231	-1.25	3.78	124	0.48	1.58	209	-0.23	5.81	84	-2.12	1.57
P2	744	0.38	1.3	100	0.75	1.07	204	0.15	0.88	142	0.52	0.67	215	0.51	2.08	83	0.17	0.26
P3	813	1	1.41	102	0.66	0.54	250	0.89	0.99	148	0.81	1.02	211	1.69	2.56	102	1.22	0.87
P4	783	0.54	0.71	88	0.56	0.33	235	0.65	0.86	154	0.64	0.59	213	0.35	0.71	93	0.4	0.49

I1, 2, 3 = first, second, third permanent incisor; C = permanent canine, P1, 2, 3, 4 = first, second, third and fourth permanent premolar; M1, 2, 3 = first, second and third molar